

INVASIVE CLEAVAGE OF NUCLEIC ACIDS

This is a Continuation-In-Part of co-pending application Serial No. 08/682,853, filed July 12, 1996, which is a Continuation-In-Part of co-pending application Serial No. 08/599,491, filed on January 24, 1996.

5 FIELD OF THE INVENTION

The present invention relates to means for the detection and characterization of nucleic acid sequences and variations in nucleic acid sequences. The present invention relates to methods for forming a nucleic acid cleavage structure on a target sequence and cleaving the nucleic acid cleavage structure in a site-specific manner. The 5' nuclease activity of a variety of enzymes is used to cleave the target-dependent cleavage structure, thereby indicating the presence of specific nucleic acid sequences or specific variations thereof. The present invention further provides novel methods and devices for the separation of nucleic acid molecules based by charge.

15 BACKGROUND OF THE INVENTION

The detection and characterization of specific nucleic acid sequences and sequence variations has been utilized to detect the presence of viral or bacterial nucleic acid sequences indicative of an infection, the presence of variants or alleles of mammalian genes associated with disease and cancers and the identification of the source of nucleic acids found in forensic samples, as well as in paternity determinations.

20 Various methods are known to the art which may be used to detect and characterize specific nucleic acid sequences and sequence variants. Nonetheless, as nucleic acid sequence data of the human genome, as well as the genomes of pathogenic organisms accumulates, the demand for fast, reliable, cost-effective and user-friendly tests for the detection of specific nucleic acid sequences continues to grow. Importantly, these tests must be able to create a detectable signal from samples which contain very few copies of the sequence of interest. The following discussion

examines two levels of nucleic acid detection assays currently in use: I. Signal Amplification Technology for detection of rare sequences; and II. Direct Detection Technology for detection of higher copy number sequences.

I. Signal Amplification Technology Methods For Amplification

The "Polymerase Chain Reaction" (PCR) comprises the first generation of methods for nucleic acid amplification. However, several other methods have been developed that employ the same basis of specificity, but create signal by different amplification mechanisms. These methods include the "Ligase Chain Reaction" (LCR), "Self-Sustained Synthetic Reaction" (3SR/NASBA), and "Q β -Replicase" (Q β).

Polymerase Chain Reaction (PCR)

The polymerase chain reaction (PCR), as described in U.S. Patent Nos. 4,683,195 and 4,683,202 to Mullis and Mullis *et al.* (the disclosures of which are hereby incorporated by reference), describe a method for increasing the concentration of a segment of target sequence in a mixture of genomic DNA without cloning or purification. This technology provides one approach to the problems of low target sequence concentration. PCR can be used to directly increase the concentration of the target to an easily detectable level. This process for amplifying the target sequence involves introducing a molar excess of two oligonucleotide primers which are complementary to their respective strands of the double-stranded target sequence to the DNA mixture containing the desired target sequence. The mixture is denatured and then allowed to hybridize. Following hybridization, the primers are extended with polymerase so as to form complementary strands. The steps of denaturation, hybridization, and polymerase extension can be repeated as often as needed, in order to obtain relatively high concentrations of a segment of the desired target sequence.

The length of the segment of the desired target sequence is determined by the relative positions of the primers with respect to each other, and, therefore, this length is a controllable parameter. Because the desired segments of the target sequence

become the dominant sequences (in terms of concentration) in the mixture, they are said to be "PCR-amplified."

Ligase Chain Reaction (LCR or LAR)

The ligase chain reaction (LCR; sometimes referred to as "Ligase Amplification Reaction" (LAR) described by Barany, Proc. Natl. Acad. Sci., 88:189 (1991); Barany, PCR Methods and Applic., 1:5 (1991); and Wu and Wallace, Genomics 4:560 (1989) has developed into a well-recognized alternative method for amplifying nucleic acids. In LCR, four oligonucleotides, two adjacent oligonucleotides which uniquely hybridize to one strand of target DNA, and a complementary set of adjacent oligonucleotides, which hybridize to the opposite strand are mixed and DNA ligase is added to the mixture. Provided that there is complete complementarity at the junction, ligase will covalently link each set of hybridized molecules. Importantly, in LCR, two probes are ligated together only when they base-pair with sequences in the target sample, without gaps or mismatches. Repeated cycles of denaturation, hybridization and ligation amplify a short segment of DNA. LCR has also been used in combination with PCR to achieve enhanced detection of single-base changes. Segev, PCT Public. No. W09001069 A1 (1990). However, because the four oligonucleotides used in this assay can pair to form two short ligatable fragments, there is the potential for the generation of target-independent background signal. The use of LCR for mutant screening is limited to the examination of specific nucleic acid positions.

Self-Sustained Synthetic Reaction (3SR/NASBA)

The self-sustained sequence replication reaction (3SR) (Guatelli *et al.*, Proc. Natl. Acad. Sci., 87:1874-1878 [1990], with an erratum at Proc. Natl. Acad. Sci., 87:7797 [1990]) is a transcription-based *in vitro* amplification system (Kwok *et al.*, Proc. Natl. Acad. Sci., 86:1173-1177 [1989]) that can exponentially amplify RNA sequences at a uniform temperature. The amplified RNA can then be utilized for mutation detection (Fahy *et al.*, PCR Meth. Appl., 1:25-33 [1991]). In this method, an oligonucleotide primer is used to add a phage RNA polymerase promoter to the 5' end